

Table 2 - Comparison of auditor's sea lice counts to industry-reported counts during Quarter 4 - 2010

2010 - Quarter4

	Number of Farms			<i>Lepeophtheirus salmonis</i>				<i>Caligus clemensi</i>			
	Auditted	Number Operating		All Motile Stages		Adult ♀ (with and without egg strings)		All Motile Stages		Chalimus Stages	
				Audit	BCSFA Database	Audit	BCSFA Database	Audit	BCSFA Database	Audit	BCSFA Database
Zone/ Subzone 2.3	2	6	Abundance for area	0.02	0.21	0.00	0.11	0.00	0.04	0.00	0.03
			Lower 95% CI		0.10	0.05		0.01		0.00	
			Upper 95% CI		0.32	0.18		0.07		0.06	
Zone/ Subzone 2.4	3	9	Abundance for area	3.71	2.59	1.97	1.21	0.06	0.09	0.37	0.38
			Lower 95% CI		1.19	0.50		0.02		0.08	
			Upper 95% CI		4.00	1.91		0.16		0.67	
Zone/ Subzone 3.1	0	2	Abundance for area	-	0.32	-	0.10	-	0.00	-	0.01
			Lower 95% CI		0.09	-0.01		0.00		0.00	
			Upper 95% CI		0.55	0.20		0.01		0.02	
Zone/ Subzone 3.2	4	11	Abundance for area	7.65	3.41	3.47	1.66	0.33	0.20	1.58	1.00
			Lower 95% CI		1.97	0.85		0.09		0.40	
			Upper 95% CI		4.86	2.47		0.31		1.59	
Zone/ Subzone 3.3	2	12	Abundance for area	2.08	1.08	0.54	0.42	0.46	0.22	1.12	0.33
			Lower 95% CI		0.56	0.20		0.07		0.15	
			Upper 95% CI		1.60	0.65		0.37		0.52	
Zone/ Subzone 3.4	2	6	Abundance for area	3.43	2.22	1.17	0.90	0.06	0.19	0.42	0.39
			Lower 95% CI		1.09	0.31		0.04		0.10	
			Upper 95% CI		3.35	1.48		0.33		0.68	
Zone/ Subzone 3.5	0	4	Abundance for area	-	3.13	-	1.63	-	0.06	-	0.80
			Lower 95% CI		0.71	0.00		-0.02		0.18	
			Upper 95% CI		5.56	3.25		0.13		1.41	